



PCT/GB 2004 / 0 0 1 1 0 6



INVESTOR IN PEOPLE

**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

REC'D 20 APR 2004

WIPO


PCT

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed


Dated

P. Mahoney
8 April 2004

Best Available Copy

Patents Form 1/77

Patents Act 1977
(Rule 16)



20MAR03 E793656-1 D10063
P01/7700 0.00-0306340.1

Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office

Cardiff Road
Newport
South Wales
NP10 8QQ

1. Your reference

386

2. Patent application number

(The Patent Office will fill in this part)

20 MAR 2003

0306340.1

3. Full name, address and postcode of the or of each applicant (*underline all surnames*)

CDT Oxford Limited
Greenwich House
Madingley Rise
Madingley Road
Cambridge CB3 0TX

Patents ADP number (*if you know it*)

If the applicant is a corporate body, give the country/state of its incorporation

United Kingdom

8541435001

4. Title of the invention

Method for the Production of Metal Complexes

5. Name of your agent (*if you have one*)

"Address for service" in the United Kingdom to which all correspondence should be sent (*including the postcode*)

IP Department
CDT Limited
Greenwich House
Madingley Rise
Madingley Road
Cambridge
CB3 0TX

Patents ADP number (*if you know it*)

8591448001

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (*if you know it*) the or each application number

Country

Priority application number
(*if you know it*)

Date of filing
(day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing
(day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (*Answer 'Yes' if:*

YES

- a) any applicant named in part 3 is not an inventor, or
 - b) there is an inventor who is not named as an applicant, or
 - c) any named applicant is a corporate body.
- See note (d))

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description 14

Claim(s) 3

Abstract -

Drawing(s) -

ph

10. If you are also filing any of the following, state how many against each item.

Priority documents -

Translations of priority documents -

Statement of inventorship and right to grant of a patent (Patents Form 7/77) -

Request for preliminary examination and search (Patents Form 9/77) 1

Request for substantive examination (Patents Form 10/77) 1

ph

Any other documents
(please specify)

11. I/We request the grant of a patent on the basis of this application.

Signature

E. Kirby

Date

17th March 2003

12. Name and daytime telephone number of person to contact in the United Kingdom E. Kirby
Tel: 01865 309610

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 08459 500505.*
- Write your answers in capital letters using black ink or you may type them.*
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.*
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.*
- Once you have filled in the form you must remember to sign and date it.*
- For details of the fee and ways to pay please contact the Patent Office.*

Method for the Production of Metal Complexes

Field of the Invention

The present invention relates to a method of producing metal complexes.

Background of the Invention

One class of opto-electrical devices attracting increasing attention is that using a semiconducting organic material for light emission (an electroluminescent device) or as the active component of a photocell or photodetector (a "photovoltaic" device). The basic structure of these devices is a semiconducting organic layer sandwiched between a cathode for injecting or accepting negative charge carriers (electrons) and an anode for injecting or accepting positive charge carriers (holes) into the organic layer.

In an organic light emitting device (OLED), electrons and holes are injected into the semiconducting organic layer where they combine to generate excitons that undergo radiative decay. Various classes of organic light emitting materials are known, in particular: polymers such as poly(p-phenylene vinylene) (as disclosed in WO 90/13148), polyfluorenes and polyphenylenes; the class of materials known as small molecule materials such as (8-hydroxyquinoline) aluminium ("Alq3") as disclosed in US 4,539,507; and the class of materials known as dendrimers as disclosed in WO 99/21935. These materials electroluminesce by radiative decay of singlet excitons (i.e. fluorescence) however spin statistics dictate that up to 75% of excitons are triplet excitons, i.e. quantum efficiency may be as low as 25% for fluorescent OLEDs – see, for example, *Chem. Phys. Lett.*, 1993, 210, 61, *Nature (London)*, 2001, 409, 494, *Synth. Met.*, 2002, 125, 55 and references therein.


Accordingly, considerable effort has been directed towards producing luminescence from triplet excitons (phosphorescence) by utilising metal complexes that enable triplet excitons to undergo radiative decay. Examples of complexes investigated for this purpose include lanthanide metal chelates [*Adv. Mater.*, 1999, 11, 1349], a platinum (II) porphyrin [*Nature (London)*, 1998, 395, 151] and iridium tris(phenylpyridine) [*Appl. Phys. Lett.*, 1999, 75, 4; *Appl. Phys. Lett.*, 2000, 77, 904]. Fuller reviews of such complexes may be found in *Pure Appl. Chem.*, 1999, 71, 2095, *Materials Science & Engineering*, R:

Reports (2002), R39(5-6), 143-222 and Polymeric Materials Science and Engineering (2000), 83, 202-203.

One approach to synthesis of these metal complexes is described in, for example, WO 02/15645, US 2002/034656, JP 2002-105055 and references therein. This is a two-step process, as illustrated in Scheme 1 for an asymmetric complex comprising phenylpyridine:

- 1) Formation of a chloro-bridged dimer (a).
- 2) Reaction of the dimer in the presence of a base and ligand to form the end product (b).

Scheme 1



L = bidentate ligand

It will be appreciated by the skilled person that this two-step process for iridium complexes is applicable to a wider range of metal complexes that form bridged dimers. Other metals that form halo-bridged dimers include rhodium, palladium and platinum (see, for example, Inorg. Chem. 2002, 41, 3055).

These prior art syntheses of organometallic complexes entail a two-step process because the chloro-bridged dimer (a) is a relatively stable material. The chloro bridge of the dimer cannot be broken by a ligand such as phenylpyridine and so the second step requires a ligand L that is capable of breaking the chloro bridge, thus making this synthesis suitable only for asymmetric complexes (i.e. those wherein L is not PPy).

It is possible to form tris(phenylpyridine) iridium (III) by reaction of phenylpyridine with dimer (a) by performing the reaction in the presence of a salt such as silver triflate as


described in Inorganic Chemistry (1994), 33(3), 545-50 and Chemical Communications (Cambridge, United Kingdom) (2001), (16), 1494-1495. This approach has the drawback that residual silver may be present in the end product which can be difficult to separate from the end product and which may be detrimental to OLED performance. Furthermore, this approach has only been reported for the synthesis of symmetric complexes (i.e. those wherein L = PPy). "Symmetric complexes" as used herein means complexes wherein all ligands attached to the metal are the same; "asymmetric complexes" as used herein means complexes wherein at least one of the ligands attached to the metal differs from at least one other ligand attached to the metal.

An alternative synthesis of iridium complexes is described, for example, in WO 02/60910 and US 2002/034656 wherein *tris*-acetylacetonate iridium(III) is reacted with 3 equivalents of phenylpyridine. This has the drawback that *tris*-acetylacetonate iridium is relatively expensive in comparison to iridium chloride (from which it is made). It is possible to prepare *tris*-acetylacetonate iridium by reaction of iridium trichloride and 2,5-pentanedione, however this adds an additional step to the synthesis (the inventors have found that it is not possible to react iridium trichloride with 2-phenylpyridine in the presence of 2,5-pentanedione because this results in degradation of the reactants). Furthermore, this one-step approach does not allow formation of asymmetric complexes.

In view of the aforementioned drawbacks of prior art syntheses, it is an object of the invention to provide a method of reacting halo-bridged iridium, rhodium, platinum or palladium dimers with ligands that are incapable of breaking the halo bridge of the dimer, but without the need for metal containing reagents other than the dimer itself. It is a further object of the invention to provide a one-pot process for synthesising said complexes that is suitable for synthesising both symmetric and asymmetric complexes (by "one-pot" as used herein is meant a process wherein the intermediate halogen bridged dimer formed in the first step is not separated from some or all of the components of the reaction mixture in which it is formed prior to the second step). It is a yet further object of the invention to provide a method for synthesising said complexes in a one-pot process without the need for costly starting materials.

Summary of the Invention

In a first aspect the invention provides a method of forming a metal complex of formula $M(\text{Ar}^1\text{Ar}^2)_n\text{L}$ comprising the step of reacting a compound of formula (I) with a bidentate ligand L:



wherein Ar^1 and Ar^2 are each independently an optionally substituted aryl or heteroaryl; $\text{Ar}^1\text{-Ar}^2$ forms at least one carbon-M bond by reaction of M with a carbanion of $\text{Ar}^1\text{-Ar}^2$; L is a compound of formula $\text{Ar}^1\text{-Ar}^2$; M is iridium, rhodium, platinum or palladium; Hal is a halogen; and n is a number from 1-3 having a value necessary to satisfy the valency of metal M,

characterised in that the reaction is performed in the presence of an enabling ligand that is capable of breaking the halogen bridge of the compound of formula (I).

By "bidentate ligand" is meant a ligand which, when present in a metal complex, is bound to the metal by two ligand-metal bonds. Each of the ligand-metal bonds may be a covalent or co-ordination bond.


Preferably, each Hal is bromine, chlorine or iodine, most preferably chlorine.

Preferably, $\text{Ar}^1\text{-Ar}^2$ is phenylpyridine.

In a first embodiment, $\text{Ar}^1\text{-Ar}^2$ and L are the same. In a second embodiment, $\text{Ar}^1\text{-Ar}^2$ and L are different.

In one embodiment of the first aspect of the invention, the enabling ligand is a monodentate ligand, more preferably a ligand selected from optionally substituted pyridine and triarylphosphine.

In another embodiment of the first aspect of the invention, the enabling ligand is a bidentate ligand of formula (IIb):



(IIb)

wherein each R is independently selected from H or a substituent. Preferably, each R is hydrogen.


Preferably, the ligand of formula (IIb) is formed by treating a corresponding protonated compound with a metal-free base.

Preferably, the reaction is performed in a protic solvent, preferably ethylene glycol.

In a preferred embodiment, the invention provides a method of forming a metal complex of formula $M(Ar^1Ar^2)_nL$ comprising a first step of preparing a compound of formula (I) by reacting a compound of formula $M(Hal)_m$ with a compound of Ar^1-Ar^2 and a second step according to the first aspect of the invention, wherein m is a number necessary to satisfy the valency of M, characterised in that the first and second steps are performed in a one-pot process.

In a second aspect, the invention provides a method of forming a metal complex comprising:

a) a first step of reacting a compound of formula $M(Hal)_m$ with a compound of formula Ar^1-Ar^2 to form a compound of formula (I):




and

b) a second step of reacting the compound of formula (I) with a reactive ligand that is capable of breaking the halogen bridge of the compound of formula (I)

wherein Ar^1 and Ar^2 are each independently an optionally substituted aryl or heteroaryl; Ar^1-Ar^2 forms at least one carbon-M bond by reaction of M with a carbanion of Ar^1-Ar^2 ; L is a compound of formula Ar^1-Ar^2 ; M is iridium, rhodium, platinum or palladium; Hal is a halogen; m is a number from 2-8 and n is a number from 1-3, m and n each having a value necessary to satisfy the valency of metal M,

characterised in that the first and second steps are performed in a one-pot process.

In a third aspect, the invention provides a method of forming a metal complex comprising the step of reacting a metal halide with a ligand of formula (IIa):



(IIa)

wherein each R is independently selected from hydrogen or a substituent. Preferably, each R is hydrogen.

characterised in that the reaction is performed in the presence of a metal-free base of sufficient strength to deprotonate the compound of formula (IIa).

By "metal halide" is meant a compound comprising at least two metal-halogen bonds, including compounds of formula $M(Hal)_x(L')_y$ wherein M is a metal, Hal is a halogen, L' is a mono- or multi-dentate ligand, x is greater than or equal to 2, y is greater than or equal to 0, and x+y is a number necessary to satisfy the valency of M. The term "metal halide"


as used in accordance with the third aspect of the invention also encompasses halide bridged dimers of metals.

The present inventors have surprisingly found that it is possible to form various metal complexes by reaction of a halo-bridged dimer with a bidentate ligand L in the presence of the aforementioned enabling ligands whereas said metal complexes do not form when the enabling ligand is absent. Thus, use of these enabling ligands allows a broader range both of products (symmetric and asymmetric) and of reaction conditions.

The formation of said metal complexes is made possible by the fact that the enabling ligand is capable of breaking the relatively stable halogen bridge of dimer (I) to form an a relatively reactive intermediate that is then capable of reacting with bidentate ligand Ar^1 - Ar^2 . This is illustrated in Scheme 2 wherein the enabling ligand is a monodentate ligand L^m (e.g. pyridine).

Furthermore, as shown in Scheme 2, the first step of the inventive process entails synthesis of the dihalo bridged dimer (I) as per the prior art. To produce the final product, prior art syntheses entail separation of the dimer (I) and subsequent reaction with bidentate ligand L. By contrast, the present inventors have surprisingly found that the final product (IV) may be formed by reacting the dimer (I) *in situ* with ligand L in the presence of monodentate ligand L^m , i.e. without having to first separate dimer (I). It will be appreciated, however, that the reaction can also be performed as a "two-pot" process, for example wherein the dimer (I) is formed in a first, protic, solvent such as ethylene glycol and subsequently reacted in a second, aprotic, solvent such as dichloromethane.

Scheme 2



Alternatively, certain bidentate ligands may be used as enabling ligands.

In addition to using ligands that are capable of breaking the halogen bridge as enabling ligands, these ligands may also be used as reactants ("reactive ligands") in a one-pot process for formation of asymmetric complexes. In this instance, the compound of formula (III) is the end product rather than an intermediate. Products of this type are known as disclosed in, for example, WO 02/15645 however prior art syntheses entail a two-pot process.

The present invention enables the preparation of symmetric or asymmetric metal complexes from a halo-bridged dimer, or in a one-pot process starting from the corresponding metal halide, without the need for silver salts as per the prior art. Furthermore, according to the third aspect of the invention, the present inventors have devised a method of preparing metal complexes with ligands of formula (IIb) which are formed by deprotonation of a compound of formula (IIa) but which are substantially free of any metal impurities. According to this method, complexes are prepared using a compound of formula (IIa) and a metal-free base. The ligand of formula (IIb) may be an enabling ligand as per the second embodiment or a reactive ligand that forms part of the end product as per the third embodiment of the invention.

Detailed Description of the Invention

Ar¹-Ar² and L may be the same or different. In the process of Scheme 2, L is added in the final step of the reaction, however L may be included in the starting mixture with Ar¹.

Ar^2 , particularly if $\text{Ar}^1\text{-Ar}^2$ and L are the same. If $\text{Ar}^1\text{-Ar}^2$ and L are different then addition of L in the final step is preferred in order to produce a single asymmetric complex.

In addition to being linked to M through a carbanion, $\text{Ar}^1\text{-Ar}^2$ in compounds of formula III may be linked to metal M by a metal-nitrogen bond.

Ar^1 and Ar^2 are preferably selected from the group consisting of optionally substituted phenyl and optionally substituted heteroaromatics comprising a 5 or 6 membered ring, either as a part of a monocyclic or fused ring system. Particularly preferred heteroaromatics ring systems are those comprising one or more oxygen, sulfur or nitrogen atoms.


L is selected from the same group of compounds as $\text{Ar}^1\text{-Ar}^2$, i.e. those compounds that are incapable of breaking the halogen bridge. For any method according to the invention, L and $\text{Ar}^1\text{-Ar}^2$ may be the same (for production of a symmetric complex) or different (for production of an asymmetric complex). L must be capable of displacing the enabling ligand.

Ligands capable of breaking the halogen bridge of the metal dimer (i.e. enabling ligands or reactive ligands) are selected from monodentate ligands or certain bidentate ligands.

Particularly preferred monodentate ligands are selected from the group consisting of optionally substituted phosphines, particularly triaryl- or trialkyl-phosphines, most preferably triphenylphosphine; optionally substituted electron rich 6-membered nitrogen containing heteroaromatics, preferably optionally substituted pyridine, diazines or triazines, most preferably pyridine.

Monodentate enabling ligands are preferred for synthesis of asymmetric complexes.

Suitable bidentate ligands for use as enabling or reactive ligands include bidentate ligands that bind to metal M through one or more oxygen atoms. One particularly preferred class of bidentate ligand are ligands of formula (IIb):




(IIb)

wherein each R is independently selected from hydrogen or a substituent. When two or more groups R are a substituent, they may together form an alicyclic or aromatic ring. Preferably, each R is hydrogen.

It will be appreciated that the ligand of formula (IIb) is formed by treatment of the corresponding protonated compound with a base capable of deprotonating said compound.


An alternative preferred enabling or reactive bidentate ligand is optionally substituted picolinate of formula (V):



(V)

Where two or more substituents are present, they may together form an alicyclic or aromatic ring.

A further preferred class of enabling or reactive bidentate ligand is that which binds to metal M through two nitrogen-metal bonds, in particular optionally substituted ligands of formula (VI):




(VI)

wherein each pyridyl ring is optionally substituted and wherein two or more of said substituents may form an alicyclic or aromatic ring.

Where the bidentate ligand derived from the compound of formula (IIb) is used, either as an enabling ligand or as a reactive ligand, all groups R are preferably H or C₁₋₁₀ alkyl, more preferably H. The compound of formula (IIb) is preferably deprotonated by a metal-free base. Particularly preferred metal-free bases include ammonium containing bases such as ammonium hydroxide or ammonium carbonate or organic bases such as trialkyl- or triaryl-amines, particularly C₁₋₁₀ alkylamines, most particularly triethylamine.

Suitable solvents for the one-pot process according to the invention are protic solvents, preferably diols, most preferably ethylene glycol.

Numerous examples of materials suitable for use as Ar¹-Ar² and L are known as disclosed in, for example, WO 02/15645. Preferably, each Ar¹-Ar² and each L comprises a nitrogen atom capable of forming a co-ordination bond with a metal. Examples of Ar¹-Ar² and L include the following:



X = O, S
Y = C, N

Each of the above groups Ar¹-Ar² may carry one or more substituents. Particularly preferred substituents include fluorine or trifluoromethyl which may be used to blue-shift the emission of the complex as disclosed in WO 02/45466, WO 02/44189, US 2002-117662 and US 2002-182441; alkyl or alkoxy groups as disclosed in JP 2002-324679; carbazole which may be used to assist hole transport to the complex when used as an emissive material as disclosed in WO 02/81448; bromine, chlorine or iodine which can serve to functionalise the ligand for attachment of further groups as disclosed in WO 02/68435 and EP 1245659; and dendrons which may be used to obtain or enhance solution processability of the metal complex as disclosed in WO 02/66552.

Examples

Example 1 - synthesis of Ir(ppy)₃ from dimer

A reaction mixture of [Ir(ppy)₂Cl]₂ (1.30 g, 1.21 mmol), 2-phenylpyridine (0.40 g, 2.58 mmol), 2,4-pentanedione (0.27 g, 2.70 mmol), triethylamine (0.27 g, 2.67 mmol) and anhydrous ethylene glycol (20 ml) was heated at reflux for 3 days in a nitrogen atmosphere. The reaction mixture was allowed to cool to room temperature. The yellow precipitate was filtered, washed with water (3 x 100 ml) followed by ethanol (3 x 100 ml) and finally dried. A yield of 1.36 g (99 %) of Ir(ppy)₃ was collected.

¹H [300 MHz, CDCl₃]: 7.88 (1H, d, Ar), 7.66 (1H, d-d, Ar), 7.58 (1H, d-t, Ar), 7.53 (1H, d-d, Ar), 6.85 (4H, m, Ar).

Example 2 – one-pot synthesis of Ir(ppy)₃ from IrCl₃

A reaction mixture of IrCl₃·3H₂O (0.20 g, 0.57 mmol), 2-phenylpyridine (0.30 g, 1.93 mmol) and anhydrous ethylene glycol (20 ml) was heated at reflux for 18 h in a nitrogen atmosphere. The initial black suspension turned to yellow precipitate, indicating formation of the chloro-bridged dimer. The mixture was allowed to cool to room temperature followed by addition of 2,4-pentanedione (0.18 g, 1.80 mmol) and triethylamine (0.18 g, 1.80 mmol). The mixture was heated at reflux for 3 days and then cooled to room temperature. The yellow ppt was filtered, washed with water (3 x 50 ml) followed by ethanol (3 x 50 ml) and finally dried. A yield of 0.22 g (60 %) Ir(ppy)₃ was collected.

¹H [300 MHz, CDCl₃]: 7.88 (1H, d, Ar), 7.66 (1H, d-d, Ar), 7.58 (1H, d-t, Ar), 7.53 (1H, d-d, Ar), 6.85 (4H, m, Ar).

Example 3 - synthesis of Ir(Brppy)₃ from dimer

A reaction mixture of [Ir(Brppy)₂Cl]₂ (2.60 g, 1.87 mmol), bromophenylpyridine (1.00 g, 4.27 mmol), 2,4-pentanedione (0.40 g, 4.00 mmol), triethylamine (0.40 g, 3.96 mmol) and anhydrous ethylene glycol (20 ml) was heated at reflux for 24 h. in a nitrogen atmosphere and then allowed to cool to room temperature. The yellow ppt was filtered, washed with water (3 x 100 ml) followed by ethanol (3 x 100 ml) and finally dried. A yield of 2.50 g (75 %) Ir(Brppy)₃ was collected.


¹H [300 MHz, dmso-d₆]: 8.22 (1H, d, Ar), 7.93 (1H, d, Ar), 7.82 (1H, d-t, Ar), 7.45 (1H, d-d, Ar), 7.17 (1H, d-t, Ar), 6.85 (1H, d-d, Ar), 6.49 (1H, d, Ar).

The method of the invention is suitable for the preparation of metal complexes for use as active components of organic optical devices such as organic electroluminescent or photovoltaic devices, in particular as phosphorescent materials in organic electroluminescent devices. However, it will be apparent that metal complexes prepared according to the invention are also suitable for other applications.

Although the present invention has been described in terms of specific exemplary embodiments, it will be appreciated that various modifications, alterations and / or combinations of features disclosed herein will be apparent to those skilled in the art without departing from the spirit and scope of the invention as set forth in the following claims.

Claims


- 1) A method of forming a metal complex of formula $M(Ar^1Ar^2)_nL$ comprising the step of reacting a compound of formula (I) with a bidentate ligand L:



wherein Ar^1 and Ar^2 are each independently an optionally substituted aryl or heteroaryl; Ar^1-Ar^2 forms at least one carbon-M bond by reaction of M with a carbanion of Ar^1-Ar^2 ; L is a compound of formula Ar^1-Ar^2 ; M is iridium, rhodium, platinum or palladium; Hal is a halogen; and n is a number from 1-3 having a value necessary to satisfy the valency of metal M,

characterised in that the reaction is performed in the presence of an enabling ligand that is capable of breaking the halogen bridge of the compound of formula (I).


- 2) A method according to claim 1 wherein Hal is bromine, chlorine or iodine, preferably chlorine.
- 3) A method according to any preceding claim wherein Ar^1-Ar^2 is phenylpyridine.
- 4) A method according to any preceding claim wherein Ar^1-Ar^2 and L are the same.
- 5) A method according to any preceding claim wherein Ar^1-Ar^2 and L are different.
- 6) A method according to any preceding claim wherein the enabling ligand is a monodentate ligand.
- 7) A method according to claim 6 wherein the monodentate ligand is selected from optionally substituted pyridine and triarylphosphine.
- 8) A method according to any one of claims 1-5 wherein the enabling ligand is a bidentate ligand of formula (IIb):



(IIb)

wherein each R is independently selected from H or a substituent.

- 9) A method according to claim 8 wherein the ligand of formula (IIb) is formed by treatment of a corresponding protonated compound with a metal-free base.
- 10) A method according to claim 8 or 9 wherein each R is hydrogen.
- 11) A method of forming a metal complex of formula $M(\text{Ar}^1\text{Ar}^2)_n\text{L}$ comprising a first step of preparing a compound of formula (I) by reacting a compound of formula $M(\text{Hal})_m$ with a compound of $\text{Ar}^1\text{-Ar}^2$ and a second step according to any preceding claim, wherein m is a number necessary to satisfy the valency of M, characterised in that the first and second steps are performed in a one-pot process.
- 12) A method according to any preceding claim wherein the reaction is performed in a protic solvent, preferably ethylene glycol.
- 13) A method of forming a metal complex comprising:
 - a) a first step of reacting a compound of formula $M(\text{Hal})_m$ with a compound of formula $\text{Ar}^1\text{-Ar}^2$ to form a compound of formula (I):




and

b) a second step of reacting the compound of formula (I) with a reactive ligand that is capable of breaking the halogen bridge of the compound of formula (I)

wherein Ar^1 and Ar^2 are each independently an optionally substituted aryl or heteroaryl; $\text{Ar}^1\text{-Ar}^2$ forms at least one carbon-M bond by reaction of M with a carbanion of $\text{Ar}^1\text{-Ar}^2$; L is a compound of formula $\text{Ar}^1\text{-Ar}^2$; M is iridium, rhodium, platinum or palladium; Hal is a halogen; m is a number from 2-8 and n is a number from 1-3, m and n each having a value necessary to satisfy the valency of metal M,

characterised in that the first and second steps are performed in a one-pot process.


- 14) A method of forming a metal complex comprising the step of reacting a metal halide with a ligand of formula (II):



(II)

wherein each R is H or a substituent,

characterised in that the reaction is performed in the presence of a metal-free base of sufficient strength to deprotonate the compound of formula (II).



This Page is inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ BLACK BORDERS
- ☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLORED OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REPERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images problems checked, please do not report the problems to the IFW Image Problem Mailbox